


IgG Fc domains that bind C1q but not effector Fc γ receptors delineate the importance of complement-mediated effector functions

Chang-Han Lee¹ , Gabrielle Romain², Wupeng Yan³ , Makiko Watanabe¹, Wissam Charab¹, Biliana Todorova^{4,5}, Jiwon Lee¹, Kendra Triplett³, Moses Donkor^{1,12}, Oana I Lungu¹, Anja Lux⁶, Nicholas Marshall^{1,12}, Margaret A Lindorfer⁷, Odile Richard-Le Goff^{4,5}, Bianca Balbino^{4,5,8}, Tae Hyun Kang¹, Hidetaka Tanno¹, George Delidakis¹, Corrine Alford⁹, Ronald P Taylor⁷ , Falk Nimmerjahn⁶, Navin Varadarajan², Pierre Bruhns^{4,5} , Yan Jessie Zhang^{3,10} & George Georgiou^{1,3,9–11}

Engineered crystallizable fragment (Fc) regions of antibody domains, which assume a unique and unprecedented asymmetric structure within the homodimeric Fc polypeptide, enable completely selective binding to the complement component C1q and activation of complement via the classical pathway without any concomitant engagement of the Fc γ receptor (Fc γ R). We used the engineered Fc domains to demonstrate *in vitro* and in mouse models that for therapeutic antibodies, complement-dependent cell-mediated cytotoxicity (CDCC) and complement-dependent cell-mediated phagocytosis (CDCP) by immunological effector molecules mediated the clearance of target cells with kinetics and efficacy comparable to those of the Fc γ R-dependent effector functions that are much better studied, while they circumvented certain adverse reactions associated with Fc γ R engagement. Collectively, our data highlight the importance of CDCC and CDCP in monoclonal-antibody function and provide an experimental approach for delineating the effect of complement-dependent effector-cell engagement in various therapeutic settings.

Therapeutic monoclonal antibodies (mAbs) ameliorate disease by two mechanisms that involve the binding and resultant modulation of the function of proteins associated with pathophysiology and the recruitment of effector mechanisms dependent on the crystallizable fragment (Fc) regions of antibody domains; these functions mediate, either directly or indirectly, the neutralization and clearance of targeted substrates, as well as the programming of adaptive immunity^{1,2}. Effector functions arise from the binding of the Fc domain of immunoglobulin G (IgG) to Fc γ receptors (Fc γ Rs) expressed on various leukocyte subsets and also from recruitment of the complement component C1q and the ensuing activation of the classical complement pathway. Human effector Fc γ Rs include, in addition to the well-characterized ‘classical’ (type I) receptors (in humans, Fc γ RI, Fc γ RII, Fc γ RIII and their isoforms), the lectin-like type II receptors (CD23 and CD209), TRIM21 and members of the FCRL family of receptors^{3,4}. The recruitment and signaling of type I receptors via immunocomplexes (ICs) are responsible for antibody-dependent cell-mediated cytotoxicity (ADCC) and antibody-dependent cell-

mediated phagocytosis (ADCP), reactions that have been established clinically to contribute to the mechanism of action of many therapeutic antibodies⁵. Alternatively, activation of the classical complement pathway leads to target-cell clearance by two distinct processes⁶: first, direct cell lysis that results from insertion of the membrane attack complex into the cell membrane (complement-dependent cytotoxicity (CDC)); and second, the deposition of opsonins, such as C3b, that are covalently bound onto the cell surface and in turn are recognized by complement receptors (CRs) on effector cells. The CRs activated by the deposited opsonins trigger complement-dependent cell-mediated cytotoxicity (CDCC) and complement-dependent cell-mediated phagocytosis (CDCP)^{6,7}. Additionally, activation of the classical pathway has been established to stimulate B cell and T cell adaptive immune responses⁸.

Determining in a quantitative way the relative roles of complement-dependent and Fc γ R-dependent effector mechanisms in mAb function is critical for the development of improved therapeutics^{9,10}. However, this has proven to be a very difficult problem to address

¹Department of Chemical Engineering, University of Texas at Austin, Austin, Texas, USA. ²Department of Chemical and Biomolecular Engineering, University of Houston, Houston, Texas, USA. ³Department of Molecular Biosciences, University of Texas at Austin, Austin, Texas, USA. ⁴Institut Pasteur, Department of Immunology, Unit of Antibodies in Therapy and Pathology, Paris, France. ⁵INSERM, U760, Paris, France. ⁶Institute of Genetics, Department of Biology, University of Erlangen-Nürnberg, Erlangen, Germany. ⁷Department of Biochemistry and Molecular Genetics, University of Virginia School of Medicine, Charlottesville, Virginia, USA. ⁸Université Pierre et Marie Curie, Paris, France. ⁹Department of Biomedical Engineering, University of Texas at Austin, Austin, Texas, USA. ¹⁰Institute for Cell and Molecular Biology, University of Texas at Austin, Austin, Texas, USA. ¹¹Center for Systems and Synthetic Biology University of Texas at Austin, Austin, Texas, USA. ¹²Present addresses: Department of Infection Diseases, MedImmune, Gaithersburg, Maryland, USA (M.D.), and Department of Biochemical Engineering and Structure, Merck, New Jersey, USA (N.M.). Correspondence should be addressed to G.G. (gg@che.utexas.edu).

Received 16 March; accepted 16 May; published online 12 June 2017; corrected online 27 June 2017 (details online); doi:10.1038/ni.3770

experimentally, as evinced by the longstanding debate about the relative importance of complement in the clearance of CD20⁺ B cells by mAbs (such as rituximab (Rituxan)) to the B cell-specific surface antigen CD20 (refs. 11,12). IgG isotypes capable of activating complement also bind to FcγRs to varying degrees, especially after the formation of highly aggregated ICs on target cells or viruses^{13,14}. As a result, it is not possible to distinguish, in the presence of serum, whether target-cell lysis by antibodies is dominated by ADCC or CDC and, similarly, whether phagocytosis is due to ADCP or CDCP. While ADCC and ADCP can be readily studied by well-established *in vitro* assays¹⁵, there is no straightforward manner with which to quantify the effect of CDCC and CDCP on target-cell clearance by mAbs. Because the C1q- and FcγR-binding sites on the Fc domain are proximal and partially overlap, amino-acid substitutions engineered to diminish the binding of FcγRs also eliminate the recruitment of C1q and vice versa^{16,17}.

Among the cell-elimination pathways triggered by the classical complement pathway, CDC activity is by far the easiest to measure and has been studied in great detail^{11,15}. In contrast, apart from the results of some very early, qualitative studies from more than 40 years ago, with polyclonal antibodies¹⁸, practically nothing is known about the kinetics and magnitude of target-cell elimination by CDCC and CDCP or their importance in mAb function. In the presence of serum, C3 fragments become deposited onto target cells as a result of activation of the classical pathway. Opsonized target cells are recognized by both CRs and FcγRs on effector cells. The different signaling pathways triggered by the activation of CRs and/or FcγRs ultimately result in killing of the target cells either through the release of cytotoxic proteins by effector cells or through phagocytosis. While synergism in the elimination of substrates when both CRs and FcγRs are activated has been inferred from some studies¹⁹, other reports have suggested antagonistic or opposing effects²⁰, and the precise role of CDCC and CDCP in the absence of confound effects due to FcγR engagement is not known.

RESULTS

Engineering of aglycosylated C1q-selective IgG1 Fc domains

To delineate in detail the role of CDCC and CDCP in target-cell clearance, among diverse effector functions (Fig. 1a), we focused on engineering C1q-selective, aglycosylated antibodies that lacked the ubiquitous Asn297 glycan in the Fc domain. In aglycosylated mAbs, glycan-mediated effects such as signaling via type II receptors are not relevant and, from a technical standpoint, very large libraries of aglycosylated IgG1 antibody variants with mutant Fc domains can be readily screened via display by *Escherichia coli* and multi-color flow cytometry to achieve a desired ligand-binding selectivity profile^{21–23}. Thus, we screened a library of approximately 1.5×10^9 *E. coli* transformant cells expressing full-length IgG1 antibodies with mutant Fc domains (Supplementary Fig. 1a,b and Supplementary Table 1) by flow cytometry via labeling with 10 nM phycoerythrin (PE)-labeled human C1q in the presence of a 100-fold excess (1 μM) of each human FcγR (FcγRI–FcγRIIIa) for counter-selection and in high-salt buffer to eliminate the nonspecific interaction of C1q with the *E. coli* surface (Supplementary Fig. 1c,d). Bacterial clones enriched after the seventh screening round were evaluated for binding to C1q-PE and lack of binding to high-avidity tetramers of streptavidin-tagged human FcγRIIIa_{V158}-PE, one of two natural variants of FcγRIIIa (Supplementary Fig. 1e–h). IgG1 antibodies with mutant Fc domains from selected clones were constructed with the antigen-binding fragment (Fab) of Rituxan (mAb to human CD20) and were expressed and purified from mammalian cells

either in glycosylated form or as aglycosylated antibodies through the introduction of a T299L substitution that disrupts the Asn297–X298–Thr299 (where ‘X’ is any amino acid) glycosylation motif in the Fc (Supplementary Fig. 2). We evaluated binding to human C1q or to the human FcγRs by enzyme-linked immunosorbent assay and surface-plasmon-resonance analysis with dimerized high-avidity glutathione S-transferase–tagged human low-affinity FcγRs to detect even very weak interactions (Table 1 and Supplementary Fig. 3a,b). Two Rituxan variants with mutant Fc domains were studied in detail; clone 801, which contains the amino-acid substitutions K320E (in the second domain of the immunoglobulin heavy-chain γ-constant region (Cγ2)) and Q386R (in Cγ3), bound to C1q selectively in both the aglycosylated form (the Fc domain of aglycosylated clone 801 (A801) constructed with the Fab arms of Rituxan (RA801)) and the glycosylated form (the Fc domain of glycosylated clone 801 (G801) constructed with the Fab arms of Rituxan (RG801)). As analyzed by surface plasmon resonance, RA801 had no detectable binding to any FcγR, whereas RG801 bound very weakly only to the high-affinity receptor FcγRI (Table 1 and Supplementary Fig. 3a,b). Rituxan clone 802 (with the amino-acid substitutions L245K, G246M, G247R and L351Q) showed selective binding to C1q only when expressed as an aglycosylated protein (RA802) (Table 1 and Supplementary Fig. 3a,b). Its glycosylated form, RG802, showed complete loss of binding to C1q and to effector FcγRs (Table 1 and Supplementary Fig. 3a,b). Both RA801 and RA802 displayed pH-dependent binding to the human neonatal Fc receptor (FcRn), with an affinity at pH 6.0 comparable to that of wild-type glycosylated IgG1 (Supplementary Fig. 3c); this suggested that the substitutions in these antibodies were unlikely to have a negative effect on recycling by FcRn or pharmacokinetics *in vivo*.

The binding of ICs onto cells represents an exquisitely sensitive assay for the detection of physiologically relevant IgG–FcγR interactions^{13,14}. To form ICs with defined stoichiometries, we used antibodies comprising the Fc domain of A801 or A802 fused to α-2-, 4-,6-trinitrophenyl (TNP) Fab arms (TNP-A801 and TNP-A802, respectively) and incubated those with bovine serum albumin (BSA) conjugated to an average of either 4 TNP molecules (TNP₄-BSA; low-avidity IC) or 32 TNP molecules (TNP₃₂-BSA; high-avidity ICs). ICs formed with wild-type antibody to TNP (anti-TNP) readily bound to Chinese hamster ovary (CHO) cells expressing each FcγR (Fig. 1b). In contrast, ICs of TNP-A801 and TNP-A802 of higher avidity (TNP₃₂-BSA) or lower avidity (TNP₄-BSA) showed no binding to FcγR-expressing CHO cells (Fig. 1b). We note that because of the higher surface hydrophobicity of TNP₃₂-BSA, when mixed with IgG, TNP₃₂-BSA ICs show greater background binding to CHO cells expressing no FcγRs than that of TNP₄-BSA ICs.

RA801 and RA802 mediated potent CDC activity with three widely used CD20⁺ human Burkitt's lymphoma cell lines (Raji, Ramos and Daudi), as well as with cells derived from patients with acute lymphocytic leukemia (Fig. 2a,b and Table 2). Complement activation of the classical pathway and opsonization of CD20⁺ cells by C3b was comparable to that achieved with Rituxan (Fig. 2c and Supplementary Fig. 4a,b). The lysis of Raji cells and Ramos cells via RA801 was characterized by EC₅₀ values (effector concentration for a half-maximum response) 2.3-fold lower and 4.5-fold lower, respectively, than those obtained with Rituxan (Fig. 2a,b and Table 2). We also assessed ofatumumab, an approved second-generation mAb to CD20 whose enhanced clinical activity in chronic lymphocytic leukemia has been established to result from improved CDC^{24–26}, and found that the greater CDC potency of RA801 compared favorably with that of ofatumumab: the EC₅₀ values for Raji cells and Ramos cells obtained with




Figure 1 Biochemical and functional properties of antibodies with engineered Fc domains that bind to C1q with exquisite selectivity. (a) Antibody-mediated FcγR mechanisms for cell killing (left: ADCC and ADCP) and complement-dependent mechanisms for cell killing (right: CDC, CDCC and CDCP). MAC, membrane attack complex. (b) Binding of low- and high-density ICs (left margin) formed by IgG1, A801 or A802 conjugated to TNP₄-BSA (top) or TNP₃₂-BSA (bottom) on CHO cells expressing human FcγRs¹⁴. Data are from one experiment representative of three experiments.

ofatumumab were 1.9-fold lower and 4.1-fold lower, respectively, than those obtained with Rituxan (Fig. 2d).

The enhanced CDC killing potency of ofatumumab is due to its binding to a membrane-proximal epitope in CD20 that more optimally engages C1q²⁷. Antibodies with the ofatumumab Fab fragments in combination with the clone 801 aglycosylated Fc domain (OA801) or clone 802 aglycosylated Fc domain (OA802) conferred on Raji cells and Ramos cells CDC of even greater potency than that conferred by ofatumumab (Fig. 2d and Supplementary Table 2).

After binding to antigens on the cell surface, IgG1 antibodies self-associate via the Fc domain to form hexamers that, in turn, optimally engage C1q²⁸. We investigated the possibility that the CDC activity of RA801 and RA802 might have been due to enhanced formation of hexamers. Certain amino-acid substitutions in the Fc region enhance the formation of hexamers (for example, IgG1 with three amino-acid substitutions: E345R, E430G and S440Y (RGY)), which results in the binding of C1q in solution and the activation of complement in the absence of antigen; this leads to the generation of the C4-complement-component cleavage product C4d²⁸. In contrast to mutant IgG1 that is prone to form hexamers, RA801 and RA802 did not elicit C4d

formation in solution or form oligomers that could be detected by size-exclusion chromatography (Fig. 2e). These results suggested that the activation of complement by RA801 and RA802 was probably not the result of an enhanced propensity to form hexamers, a conclusion also supported by the structural analysis reported below.

Analysis of RA801- and RA802-mediated CDCC

In standard ADCC assays performed with culture medium in the absence of serum, RA801 and RA802 were completely unable to lyse CD20⁺ cells (Raji or Ramos) with either peripheral blood mononuclear cells (PBMCs) or polymorphonuclear cells (PMNs) as effector cells (Fig. 3a, Table 2 and Supplementary Fig. 4c). In contrast, in the presence of serum (25% of the total volume) that had been depleted of complement component C9 (to abolish the formation of membrane attack complexes and thus cell killing via CDC), RA801 and RA802 were very efficient in mediating target-cell lysis, with EC₅₀ values slightly higher than those obtained with Rituxan under these conditions²⁹ (Fig. 3b). In these experiments, cell lysis was strictly dependent on the presence of effector cells (PBMCs or PMNs) and was solely the outcome of CDCC, given that RA801 and RA802 were unable to bind to FcγRs

Table 1 Binding properties of the engineered antibody variants

	N-gly	C1q	FcγRI	FcγRIIa _{H131}	FcγRIIa _{R131}	FcγRIIb	FcγRIIIa _{V158}	FcγRIIIa _{F158}	FcRn
IgG1	Yes	23.0	1.5	120	310	1300	195	390	623
IgG 801	No (RA801)	0.108 (213-fold)	NB	NB	NB	NB	NB	NB	959
	Yes (RG801)	0.385 (60-fold)	648 (0.002 fold)	NB	NB	NB	NB	NB	283.
IgG 802	No (RA802)	0.145 (158-fold)	NB	NB	NB	NB	NB	NB	110
	Yes (RG802)	NB	NB	NB	NB	NB	NB	NB	289.

Apparent dissociation constant (K_D) values (nM) for the binding of RA801 and RA802 (far left) to human C1q, human FcγRs (two natural variants of FcγRIIa (His131 and Arg131) and FcγRIIIa (Phe158 and Val158), with variant amino acids subscripted) or human FcRn (assessed at a pH of 6.0) (top row), estimated with a global two-state binding model for C1q binding, the langmuir 1:1 binding model for the binding of the high-affinity receptor FcγRI or a bivalent binding model for dimeric forms of the low-affinity FcγRs, as described²²; results were calculated by the following equation: K_D (glycosylated IgG)/ K_D (IgG 801 or 802). NB, no binding detected. Data are representative of three experiments.




Figure 2 *In vitro* complement activation by C1q-specific antibodies to CD20. (a,d) Lysis (by CDC) of CD20⁺ Raji cells (left) or Ramos cells (right) opsonized by various concentrations (horizontal axes) of Rituxan or its variants RA801 or RA802 (a) or ofatumumab or its variants OA801 or OA802 (d), or isotype-matched control antibody (Isotype), plus 25% pooled human serum, assessed 1 h after the addition of cells and antibody. (b) Lysis of patient-derived acute lymphocytic leukemia cells (ALL) opsonized by antibodies as in a (key), assessed as in a,d. (c) Deposition of C3b on CD20⁺ Daudi (left) or DB (right) human B cell lymphoma cells opsonized by culture for 15 or 30 min (horizontal axis) with antibodies as in a (key); results are presented relative to those of cells opsonized with isotype-matched control antibody (clinical-grade trastuzumab). (e) Generation of C4d by serum alone, heat-aggregated IgG (HAG; positive control) or antibodies (as in a) in solution (key); inset, size-exclusion fast protein liquid chromatography of Rituxan, RA801 and RA802 (key). Data are from one experiment representative of three experiments (error bars, s.d.).

and did not exhibit any CDC activity in serum depleted of C9 or any CDCC activity in in serum depleted of C1q (Fig. 3c,d). Thus, in serum, the efficacy of the lysis of CD20⁺ cells by RA801 and RA802, which perform only CDCC, was at least comparable to or greater than that of Rituxan, even though Rituxan executed target-cell killing via both FcγR-mediated mechanisms (ADCC) and CR-dependent mechanisms (CDCC). CR3 is a key receptor for the lysis of complement-opsonized target cells by effector cells^{26,30}. The addition of blocking antibody to CR3 inhibited the lysis, by PBMCs and PMNs (as effector cells), of RA801-opsonized Raji cells in serum depleted of C9 (Fig. 3e and Supplementary Fig. 4e). Antibody blockade of CR4 also inhibited the lysis of RA801-opsonized cells by PBMCs and PMNs^{26,30} (Fig. 3e and Supplementary Fig. 4e). Thus, CR3 and CR4, which are expressed by a variety of myeloid cells, have an important role in CDCC. Finally, in assays performed in complete blood, a condition more relevant to *in vivo* settings, RA801 and RA802 were slightly more effective than Rituxan at achieving lysis of Raji cells and Ramos cells

(Fig. 3f,g, Table 2 and Supplementary Fig. 4f). We concluded that at least within the time frame and conditions used in these experiments, complement activation made a major, if not dominant, contribution to the elimination of antibody-opsonized CD20⁺ cells.

Quantitative analysis of NK cell-mediated CDCC

Natural killer (NK) cells are by far the most effective cytolytic leukocyte population among PBMCs *in vitro*. NK cells can form stable conjugates with antibody-opsonized target cells via either FcγRIIIa (CD16a) or CRs^{26,30–32}; this results in the release of cytotoxic granules, which leads to apoptosis and target-cell lysis. We used high-throughput single-cell killing assays to evaluate, at high resolution, details of the kinetics of the killing of Raji cells by human NK cells. We used single-cell assays because the time constants for processes such as the formation of synapses between effector and target cells, time required for cell lysis and the proficiency with which effector cells perform multiple killings cannot be determined by conventional macroscopic

Table 2 *In vitro* potency of the engineered antibody variants

Condition	PHS Effector	CDC		ADCC		ADCP	CDCC (or ADCC + CDCC)		CDCP	CMC (or ADCC + CMC)	
		+C9	–	–	–	–	–C9	–	–	+C9	–
		None	PBMC	PMN	PMN	MΦ	PBMC	PMN	MΦ	PBMC	PMN
EC ₅₀ Raji	Isotype	No response									
	Rituxan	8.4 ± 1.7	9.5 ± 0.4	11.6 ± 0.8	0.04 ± 0.005		2.5 ± 0.1	1.9 ± 0.1	17.4 ± 0.2	1.0 ± 0.1	0.8 ± 0.02
	RA801	3.7 ± 0.4	–	–	–		2.6 ± 0.2	2.1 ± 0.2	3.7 ± 0.04	0.5 ± 0.02	0.4 ± 0.02
	RA802	5.0 ± 1.0	–	–	–		6.2 ± 0.3	4.6 ± 0.2	4.6 ± 0.1	0.7 ± 0.03	0.6 ± 0.05
EC ₅₀ Ramos	Isotype	No response									
	Rituxan	0.7 ± 0.1	0.8 ± 0.06	2.5 ± 0.2	NA		0.4 ± 0.03	0.3 ± 0.02	NA	0.2 ± 0.01	0.1 ± 0.01
	RA801	0.2 ± 0.02	–	–	NA		0.3 ± 0.02	0.2 ± 0.01	NA	0.03 ± 0.002	0.02 ± 0.001
	RA802	0.4 ± 0.02	–	–	NA		0.5 ± 0.03	0.4 ± 0.02	NA	0.07 ± 0.005	0.06 ± 0.004

Lysis of antibody-opsonized Raji and Ramos cells (antibodies, second column) by various effector cells (third row), in the absence of serum (–) or in the presence of pooled human serum left undepleted (+C9) or depleted of C9 (–C9) (second row); results are presented as EC₅₀ values (nM) and are 'stratified' by effector function (above table). MΦ, macrophage; NA, not assayed; CMC, complement-mediated cytotoxicity (including CDC and CDCC). Data are representative of three experiments (average ± s.d.).




Figure 3 Killing of CD20⁺ cells by CDCC. (a,b) Lysis of Raji tumor cells (opsonized by various concentrations (horizontal axes) of isotype-matched control antibody (as in Fig. 1d), Rituxan, RA801 or RA802 (key)) by PBMCs (as effector cells), at an effector cell/tumor cell ratio of 10:1, in RPMI-1640 medium without pooled human serum (No PHS) (a) or supplemented with 25% pooled human serum depleted of C9 (PHS – C9) (b); results are presented relative to those obtained by incubation in SDS lysis buffer. (c) Lysis (by CDC) of Raji cells opsonized by various concentrations (horizontal axes) of IgG antibodies as in a (key), assessed in the presence of undepleted serum (+C9) or serum depleted of C9 (–C9). (d) Lysis of Raji cells (opsonized by 20 μg/ml, 4 μg/ml or 0.8 μg/ml (dark to light bar shading) of isotype-matched control antibody, Rituxan or RA801 (key)) by PBMCs or PMNs in RPMI-1640 medium supplemented with 25% serum depleted of C1q. (e) Lysis of Raji cells (opsonized by 4 μg/ml, 0.8 μg/ml or 0.16 μg/ml (shading as in d) of antibodies as in d (key)) by CDCC via PBMCs or PMNs (above plots) coated with no antibody (–) or 10 μg/ml of anti-CR3 (α-CR3) or anti-CR4 (α-CR4) (below plots), in RPMI-1640 medium supplemented with 25% serum depleted of C9. (f) Lysis of Raji cells (opsonized as in a) by PBMCs, in RPMI-1640 medium supplemented with 25% pooled human serum. (g) Lysis of Raji cells (opsonized with 10 μg/ml, 2.5 μg/ml or 0.6 μg/ml (shading as in d) of antibodies as in a (key)) by PBMCs, in RPMI-1640 medium supplemented with 25% pooled whole blood. Data are from one experiment representative of three experiments (a–d,f,g) or two experiments (e) (error bars, s.d.).

assays. In brief, fluorescence-labeled NK cells from two healthy donors were incubated in nanowells together with an average of one to three Raji (target) cells that had been opsonized with either Rituxan or RA801. Cell motility and lysis were monitored by time-lapse imaging microscopy in nanowell grids (TIMING)³³ (Supplementary Fig. 5). The time to establish stable conjugation, the duration of conjugation before tumor-cell apoptosis and the time to tumor-cell apoptosis were determined for >150 single-cell killing events per sample (Fig. 4a,b). We observed no significant difference, following opsonization with Rituxan or RA801, in the frequency of one- or two-target-cell killing events (Fig. 4b). Mechanistically, killing via RA801 involves the formation of synapses between CRs and opsonins deposited on target cells, whereas with Rituxan, synapses can form both as a result of opsonin deposition and via the binding of antibody to FcγRIIIa on NK cells. The similar overall NK cell-killing activity or serial killing of more than one cell induced by the two mAbs indicated that either CR engagement had a dominant role in cytotoxic synapse formation or that, alternatively, in the presence of serum, FcγRIIIa-mediated cell killing was suppressed. Consistent with the latter hypothesis, complement activation has been shown to inhibit FcγRIIIa-dependent ADCC²⁰.

Notably, for both donors noted above and at an effector cell/target cell ratio of either 1:1 or 1:2, the times noted above (for establishment of conjugation, duration of conjugation before apoptosis and tumor-cell apoptosis) were all shorter for cells opsonized with RA801 than for those opsonized with Rituxan (Fig. 4c–e). Additionally, by quantifying the displacement of cells while out of or in contact with the target cells, we found that NK cells displayed greater motility

in the presence of RA801-coated Raji cells than in the presence of Rituxan-coated Raji cells (Fig. 4f). This effect might have been related to the somewhat more efficient complement activation by RA801 and the ensuing deposition of inactivated C3b, whose receptor, CR3, is expressed on human NK cells^{26,30–32}. Collectively, our data revealed that CDCC could be a major mechanism in the cytotoxic effect of NK cells on antibody-opsonized target cells.

Analysis of CDCC by RA801 and RA802

In the absence of serum, RA801- or RA802-opsonized CD20⁺ cells were not phagocytosed by classically activated (M1) macrophages, even at the highest concentration of antibody tested (400 nM) (Fig. 5a). However, RA801 and RA802 were effective in performing phagocytosis by macrophages via complement when human serum depleted of C9 was added to the assay (Fig. 5b,c). The finding that phagocytosis with RA801 or RA802 occurred only in the presence of serum unequivocally demonstrated that it was mediated by complement deposition; i.e., it was due solely to CDCC. In the presence of serum, phagocytosis of RA801- or RA802-opsonized cells was slightly more effective than that of Rituxan-opsonized cells, with EC₅₀ values of 3.7 ± 0.04 nM, 4.6 ± 0.1 nM and 17.4 ± 0.2 nM (mean ± s.d.), respectively (Fig. 5b,c). Notably, at saturating concentrations of antibody, the presence of serum markedly increased the magnitude of phagocytosis and resulted in >80% clearance of Raji cells with either RA801 or Rituxan, compared with 30% cell clearance observed with ADCC (Fig. 5b,c). However, consistent with published reports³⁴, serum increased the EC₅₀ for phagocytosis via Rituxan (17.4 ± 0.2 nM with serum, compared with 37 ± 5 pM for medium only (mean ± s.d.); Fig. 5b,c).




Figure 4 Quantitative analysis of RA801-mediated CDCC. (a) Features of single-cell analysis of target-cell killing by TIMING³³. AnnV, annexin V. (b) Frequency of the killing of Raji cells (target cells opsonized by Rituxan or RA801 (key)) by human NK cells (effector cells from two donors; far left), at various effector cell/target cell (E:T) ratios (left margin), in the presence of 25% serum depleted of C9, monitored by TIMING and presented as 0, 1 or 2 target cells killed (bar shading; middle of plot); numbers in bars indicate percent for each bar section. (c–e) Kaplan-Meier curves of the time to establish stable conjugation (t_{seek}) (c), total duration of conjugation before tumor-cell apoptosis ($t_{contact}$) (d) and time between first contact and tumor-cell apoptosis (t_{death}) (e) for NK cells (from two donors; key) incubated with Raji tumor cells (at a ratio of 1:1) opsonized with Rituxan or RA801 (key). (f) Motility of NK cells (donor 1, >840 cells; donor 2, >400 cells) in the presence of Raji cells opsonized with Rituxan or RA801 (key), with cells at a ratio of 1:1, presented as displacement (d_{well}) between two time-lapse images (displacement out of contact and in contact with the target cell computed separately³³). Each symbol represents an individual cell; small horizontal lines indicate the average (\pm s.d.). * $P \leq 0.05$, ** $P \leq 0.0001$ and *** $P \leq 0.00001$ (log-rank test (c–e) or one-way analysis of variance test (f)). Data are from two independent experiments ($n \geq 350$ cells).

Anti-tumor effects by complement in an *in vivo* model

We first established that RA801 did not bind to mouse FcγRs but was able to activate mouse complement and mediate CDC (Supplementary Fig. 6a–c). We evaluated the anti-tumor effect of RA801 in the treatment of established subcutaneous tumors formed by Ramos cells in outbred nude mice, which lack a thymus and mature T cells but produce macrophages and NK cells. When the tumors reached a diameter of 40–45 mm², the mice were treated three times with mAb (10 mg per kg body weight), administered intraperitoneally. Nearly complete inhibition of tumor growth was observed with both Rituxan and RA801, despite the fact RA801 could not bind to FcγRs or perform ADCC or ADCP (Fig. 6a).

The clearance of EL4 mouse lymphoma cells expressing human CD20 (EL4-hCD20 cells) by Rituxan has been reported by some investigators to be completely dependent on FcγRs³⁵, while others have found that either complement alone or both complement and FcγRs are important, in a manner that depends on tumor-cell burden and location²⁶. We implanted 1×10^5 luciferase-expressing EL4-hCD20 cells subcutaneously into nude mice of the NMRI strain that were deficient in the transcription factor Foxn1 (NMRI-Foxn1^{-/-} mice) and then treated the mice three times with RA801 or Rituxan (5 mg per kg body weight), administered intravenously. No significant difference between Rituxan-treated mice and RA801-treated mice was observed in terms of tumor growth, with each antibody being more effective than vehicle alone (Fig. 6b,d). In NMRI-Foxn1^{-/-} mice lacking genes encoding all effector IgG receptors (*Fcgr1*, *Fcgr2b*, *Fcgr3* and *Fcgr4*; called 'FcγR-null' here), the anti-tumor effect of Rituxan was abolished (Fig. 6c,d).

Rituxan has weak complement activity with mouse serum^{36,37}, and in mice in which FcγR-dependent effector mechanisms were abolished, complement activation alone was not sufficient to retard tumor growth (Fig. 6c,d). In contrast, administration of RA801 had a significant effect in slowing tumor growth in FcγR-null mice (Fig. 6c,d).

The presence of micro-aggregates in antibody preparations can trigger systemic, FcγR-mediated anaphylactic responses. It has been reported that intravenous administration of heat-aggregated Rituxan results in a rapid reduction in body temperature³⁸. In contrast, we observed no change in body temperature in mice given injection of heat-aggregated RA801 (Fig. 6e). Separate from that, the engagement of FcγRs by anti-CD20 has been shown to lead to decreased surface expression of CD20 on B cells, a process that is mediated in part by the co-ligation of FcγRIIb and internalization of antibody-antigen complexes *in cis*^{39,40}. Consistent with those observations, Raji cells incubated with Rituxan showed a significant decrease in surface expression of CD20, with a reduction of >40% by 6 h (Fig. 6f and Supplementary Fig. 6d). In contrast, cells incubated with RA801 showed no decreasing expression of surface CD20 (Fig. 6f and Supplementary Fig. 6d). Collectively, these findings revealed that mAbs that activated only complement had a strong anti-tumor effect in mouse models and, in addition, were able to circumvent detrimental effects of IgG1 antibodies known to be dependent on FcγR binding.

Structural analysis of the clone 801 Fc

To understand the molecular basis of the unique selectivity of the clone 801 Fc fragment for C1q, we solved the crystal structures of the




Figure 5 CDCP of CD20⁺ cells. (a,b) Phagocytosis of antibody-opsonized Raji cells (antibodies in key; concentration, horizontal axes) by monocyte-derived human M1-macrophages in RPMI-1640 medium without serum (a) or with serum depleted of C9 (b). (c) Fluorescence microscopy analyzing the phagocytosis of Raji cells, stained with the cell-permeant dye calcein-AM (green) by M1 macrophages, stained with anti-CD14 and anti-CD11b with allophycocyanin (red), showing Raji cells without macrophages (top left) or macrophages without Raji cells (bottom left), with opsonization by isotype-matched control antibody, Rituxan, RA801 or RA802 in the presence of serum depleted of C9 (top row, low magnification; bottom row, high magnification); top and right 'strips' show different views (xz and yz) of the same cells at a confocal plane. Scale bars, 20 µm. Data are from one experiment representative of three experiments (error bars (a,b), s.d. of technical triplicates).

Fc domain of both the aglycosylated clone 801 (A801-Fc) and glycosylated clone 801 (G801-Fc) at 2.3 Å and 3.2 Å, respectively (Table 3). The most striking feature of the A801-Fc structure was the extreme asymmetric orientation of C γ 2 in chain B (C γ 2_B), relative to that of C γ 3 (Fig. 7a). Topologically, the orientation of C γ 2 relative to C γ 3 is described by the C γ 2–C γ 3 dihedral angle, which is defined from the C α atoms of Tyr300 and Tyr319 in C γ 2, and Gln362 and Met428 in C γ 3 (ref. 41). Chain A of A801-Fc had a dihedral angle of –20.5°, which was slightly smaller than yet comparable to the dihedral angles of chain A observed in the ten glycosylated Fc (G-Fc) structures (–27° ± 4.1° (mean ± s.d.)) or the comparable angle (–25° ± 9.4°) in the four aglycosylated or deglycosylated human Fc proteins (A-Fc) in the Protein Data Bank (PDB) (Fig. 7a and Supplementary Table 3). However, chain B of A801-Fc showed an extreme twist and packed against C γ 2_A, which resulted in a dihedral angle of 136° (Fig. 7a). As a comparison, in all other Fc domains in the PDB, chains A and B have similar dihedral angles (–23° ± 3.8° for ten G-Fc structures, and –25° ± 1.9° for four A-Fc structures). This extreme twist was observed only for A801-Fc, but G801-Fc showed similar dihedral angles for both chain A and chain B, comparable to those in other Fc structures (Supplementary Table 3). Although it is well established that the two homodimeric Fc chains crystallize with distinct temperature factors, dynamics and geometries for each chain⁴¹, the different orientation of C γ 2_A and C γ 2_B relative to that of C γ 3 in A801-Fc is unprecedented, to our knowledge. No direct physical contact of C γ 2 with other molecules in the asymmetric unit was observed (Supplementary Fig. 7a), which suggested that the dihedral angle in chain B of A801-Fc was probably not the result of crystal-packing artifacts. We used cross-linking




Figure 6 *In vivo* activity of RA801. (a) Growth of Ramos cell xenografts (1×10^6 cells) implanted subcutaneously into nude mice ($n = 9$) treated (downward arrows) with PBS or 200 µg of Rituxan or RA801 (key) administered after palpable tumors had developed, at day 7, 10 and 13 after tumor implantation. $^{**}P < 0.0001$, versus PBS treatment (Student's *t*-test). (b,c) Growth of luciferase-expressing EL4-hCD20 cell xenografts (1×10^5 cells) in NMRI-*Foxn1*^{–/–} nude mice (Fc γ R-WT; $n = 10$ per group) (b) or NMRI-*Foxn1*^{–/–} Fc γ R-null mice ($n = 10$ per group) (c) treated (upward arrows) with PBS or 100 µg of Rituxan or RA801 (key) administered on day 1, 2 and 3 after implantation; results are presented as tumor flux over time. $^{*}P < 0.01$, versus PBS treatment (Mann-Whitney test). (d) Pseudocolor images of tumors from NMRI-*Foxn1*^{–/–} nude mice (Fc γ R-WT) and NMRI-*Foxn1*^{–/–} Fc γ R-null nude mice (left margin) as in b,c, showing tumor intensity (high (red) to low (blue)) at day 10 after implantation. (e) Change in body temperature (Δ temp) in C57BL/6J mice ($n = 3$) at various times (horizontal axis) after intraperitoneal injection of 600 µg of heat-aggregated mouse Fab, IgG2a-Rituximab or RA801 (key). (f) Internalization of IgG into CD20⁺Fc γ RIIb⁺ Raji cells incubated for 0, 2, 4 or 6 h (horizontal axis) with 100 nM isotype-matched control antibody, Rituxan, RA801 or RA802 (key), assessed by flow cytometry and presented as the mean fluorescence intensity (MFI) of surface-bound IgG. Data are from one experiment representative of three independent experiments (a,e,f; error bars, s.d.) or are from two independent experiments (b–d; error bars (b,c), s.d.).

with the carbodiimide EDC and liquid chromatography–tandem mass spectrometry to determine whether the observed crystal structure reflected the conformation of A801-Fc in solution. We observed a cross-linked product in A801-Fc between a peptide with Lys334 in chain A and a second peptide with Glu269 in chain B (Fig. 7b). In the highly asymmetric A801-Fc structure, the side chains of Lys334 in chain A and Glu269 in chain B were only 3.6 Å apart (Fig. 7b), a distance compatible with the formation of cross-linkage via EDC (~4 Å). In contrast, Lys334 in chain A and Glu269 in chain B were 24.1 Å apart in G-Fc (PDB accession code 3AVE) and were 24.1 Å apart in A-Fc (PDB accession code 3S7G) (Fig. 7b). Thus, the results of the crosslinking experiments (Fig. 7c) were fully consistent with, and provided independent support for, the proposal that the crystal structure of A801-Fc reflected the conformation of the molecule in solution.

In the A801-Fc structure, loops BC and FG in C γ 2_B formed extensive interactions with β -strands (A–B) in C γ 2_A and buried 404 Å² at

Table 3 Data collection and refinement statistics for G801-Fc and A801-Fc

Data collection	G801-Fc	A801-Fc
Space group	P 1 2 ₁ 1	P 6 ₅
Cell dimensions		
a, b, c (Å)	93.6, 141.0, 98.9	66.3, 66.3, 370.4
α, β, γ (°)	90.0, 117.7, 90.0	90.0, 90.0, 120.0
Resolution (Å)	40.0–3.22 (3.28–3.22)	50.0–2.25 (2.29–2.25)
R _{sym} or R _{merge}	0.150 (0.808)	0.089 (0.826)
I/σI	9.69 (1.55)	19.27 (1.24)
Completeness (%)	99.90 (99.80)	99.30 (96.40)
Redundancy	3.6 (3.6)	5.1 (3.7)
Refinement		
Resolution (Å)	40.0–3.22 (3.28–3.22)	50.0–2.25 (2.29–2.25)
No. reflections	36958	81432
R _{work} /R _{free}	0.252/0.292	0.197/0.243
No. atoms	11844	6713
Protein	11351	6588
Sugar atoms	493	0
Water	0	125
B-factors (Å ²)		
Protein	84	64
Sugar atoms	101	NA
Water	NA	51
r.m.s. deviations		
Bond lengths (Å)	0.002	0.005
Bond angles (°)	0.61	0.82

Shell of highest resolution is in parenthesis. NA, not applicable.

the interaction interface⁴² (Fig. 7d). The principal contacts at this surface were dominated by hydrophobic interactions between residues Phe241, Phe243 and Val264 of chain A with backbone atoms of residues Glu269, Asp270, Lys326 and Pro329 in chain B. Furthermore, a salt bridge was formed between Asp265 of chain A and Lys326 of chain B.

The A801-Fc structure suggested a molecular basis for the selective binding to C1q and lack of binding to FcγRs. The K320E amino-acid substitution lay within the predicted C1q-binding region^{43–46} and probably mediated binding to C1q by electrostatic interactions. The binding of A801-Fc to C1q was probably the result of more subtle effects probably related to differences in the conformational flexibility of Cγ2. Within the A801-Fc structure, the apex of Cγ2_B folded over to support and stabilize Cγ2_A at the region normally occupied by the glycan (Supplementary Fig. 7b). The B factor of Cγ2_A was substantially lower than that of Cγ2_B (63 Å² versus 93 Å²; Supplementary Fig. 7b). By comparison, Cγ2 in A-Fc structures²³, which showed no binding to C1q, exhibited high B factors for both chain A and chain B (for example, 98 Å² and 98 Å², respectively, in PDB accession code 3S7G), which was much larger than the B factors of Cγ3 in the respective structures (Supplementary Fig. 7b). Other biochemical data has also shown that removal of the Asn297 glycan increases the conformational flexibility of Cγ2 (ref. 23). We postulate that the enthalpic contribution from the K320E substitution, probably coupled with the lower flexibility of Cγ2_A, was a key reason for the efficient binding of A801-Fc to C1q.

X-ray structures also provided an explanation for the complete loss of binding of FcγR to A801-Fc. With IgG1, FcγRs dock on both Cγ2_A and Cγ2_B and interact with two subsites centered near the C'E loop of one chain and the hinge LLPP motif of the other⁴⁷. Interactions with both subsites are needed for FcγR binding, which results in an interaction surface of ~1,000 Å², of which about 600 Å² of interface is formed with the LLPP motif⁴⁸. However, the change in the orientation of Cγ2_B in A801-Fc precluded the simultaneous interaction of FcγRs with both subsites (Supplementary Fig. 7c).




Figure 7 Structural features of A801-Fc and G801-Fc. (a) Dihedral angles of Cγ2–Cγ3 in G-Fc (PDB accession code 3AVE; left; orange), A801-Fc (PDB accession code 5V43; middle; teal) and G801 Fc (PDB accession code 5V4E; right; purple). (b) Distance from Glu269 in chain B (E269) to Lys334 in chain A (K334) in the crystal structures of A801-Fc (PDB accession code 5V43), G-Fc (PDB accession code 3AVE) and A-Fc (PDB accession code 3S7G). (c) Tandem mass spectrometry of tryptic fragments of A801-Fc after crosslinking with EDC. (d) The chain A (teal)–chain B (gray) interface of A801-Fc. (e) Overlaid Cγ2 domain and the C'E loop (arrow) of A801-Fc (teal) and G-Fc (orange) structures; white, glycan in PDB accession code 3AVE. Data are representative of three independent experiments.

The two FcγR-binding subsites of G801-Fc were sterically accessible; however, they seemed to be highly flexible and/or disordered. Specifically, in three of the four molecules within each asymmetric unit, either the LLPP motif or the C'E loop was disordered (Fig. 7e and Supplementary Fig. 7d,e). In the only molecule within the asymmetric unit in which these regions were sufficiently ordered to be traced, the C'E loop assumed a 'soft' configuration observed only in A-Fc, which does not bind to FcγRs^{23,47} (Fig. 7e and Supplementary Fig. 7d,e). The high flexibility of the C'E loop of this 'soft' conformation would probably 'penalize' FcγR binding with a high entropic cost. Collectively, the X-ray crystal structures of A801-Fc and G801-Fc provided a structural explanation for the enhanced C1q binding and the lack of recognition by FcγRs.

DISCUSSION

Effector cells express both FcγRs and CRs; therefore, in the presence of serum, ADCC (or ADCP) and CDCC (or CDCP) occur at the same time^{13–15}. Whereas ADCC and ADCP can be assayed readily without CDCC or CDCP taking place, simply through the use of medium depleted of serum and C1q, the reverse is not the case. For analysis of CDCC or CDCP alone, without ADCC or ADCP also taking place, it is necessary to block the binding of ICs to the multitude

of FcγRs expressed by effector cells. We reasoned that a much more experimentally tractable approach for the study of CDCC and CDCP would be to engineer human IgG1 Fc domains that bound only to C1q and activated complement without engaging FcγRs. For this purpose, it was critical to 'silence' FcγR binding, since even very low-affinity interactions between IgGs and FcγRs can trigger effector phenotypes^{13,14}. The engineering of Fc domains with absolute or nearly absolute C1q selectivity involved starting with aglycosylated antibodies in which binding to both C1q and FcγRs was substantially, albeit not completely, attenuated, followed by screening of very large mutant libraries for both binding to fluorescence-labeled C1q and the simultaneous absence of binding to high-avidity FcγRs. Using this approach, we isolated A801-Fc; of its two amino-acid substitutions, K320E in Cγ2 conferred substantial affinity for C1q, and Q386R in Cγ3 resulted in a radical conformational change that abolished any binding to FcγRs.

Overwhelmingly, the effector function of therapeutic antibodies is thought to be dominated by FcγR-dependent processes and/or by CDC. We demonstrated in various assays that effector-cell-mediated substrate-clearance mechanisms triggered by activation of the classical complement pathway resulted in very extensive and rapid target-cell lysis and phagocytosis. Our results suggest that the extent and kinetics of the clearance of CD20⁺ cells via CDCC or CDCP are at least comparable to those due to ADCC or ADCP, respectively. Consequently, CDCC and CDCP deserve careful consideration and optimization in studies of the mechanism of action of mAbs.

The relative importance of CDCC and CDCP probably depends heavily on the target and disease setting. First, optimal activation of complement is critically dependent on the density and organization of antigens on the surface of target cells. Second, many pathogenic cells have high expression of complement-inhibitory proteins, and clearly, in these cases, activation of the classical pathway is compromised¹¹. Third, it is well established that *in vivo* extensive activation of complement due to a considerable burden of target cells and antibody can rapidly lead to the depletion of either C1q or downstream complement components, such as C2 (ref. 49). Clearly, in such circumstances, ADCC and ADCP are probably the dominant antibody effector mechanisms. In conclusion, mAbs with absolute C1q-binding selectivity, as described here, might be useful for the eradication of pathogenic cells in certain disease states while circumventing FcγR-dependent adverse effects, as well as the lower responsiveness to mAbs in patients with low-affinity FcγR polymorphisms⁵⁰.

METHODS

Methods, including statements of data availability and any associated accession codes and references, are available in the [online version of the paper](#).

Note: Any Supplementary Information and Source Data files are available in the online version of the paper.

ACKNOWLEDGMENTS

We thank Y. Tanno for assistance with protein expression; A. Bui for assistance with liquid chromatography–tandem mass spectrometry; P. Tucker (University of Texas at Austin) for cancer cell lines; D. Lee (MD Anderson Cancer Center) for patient-derived primary acute lymphocytic leukemia cells; the Macromolecular Crystallography Facility of the University of Texas at Austin; the Berkeley Center for Structural Biology; the Advanced Light Source (supported by the Director, Office of Science, Office of Basic Energy Sciences, of the US Department of Energy under contract DE-AC02-05CH11231); the Proteomics Facility at the University of Texas at Austin (supported by grant RP110782 from the Cancer Prevention Research Training Program); and A. Nicola and the Plate-Forme d'Imagerie Dynamique (Institut Pasteur, Paris) for help with the bioluminescence experiments. Supported by the Clayton Foundation, the Institut Pasteur

(P.B. laboratory), the Institut National de la Santé et de la Recherche Médicale (P.B. laboratory), the European Research Council Seventh Framework Program (ERC-2013-CoG 616050 for the P.B. laboratory), the Pasteur–Paris University International PhD program (B.B.), the Cancer Prevention Research Training Program (RP140108 to M.D.; RP160015 to H.T.; and RP130570 to N.V.), the American Cancer Society (123506-PF-13-354-01-CDD to N.M.), Uehara Memorial Foundation (H.T.), Japan Society for the Promotion of Science (H.T.), Deutsche Forschungsgemeinschaft (CRC1181-A07 to F.N.), the US National Institutes of Health (R01CA174385 to N.V.; and R01 GM104896 to Y.J.Z.) and the Welch Foundation (F-1778 to Y.J.Z.). The content is solely the responsibility of the authors and does not necessarily represent the official views of the Cancer Prevention and Research Institute of Texas.

AUTHOR CONTRIBUTIONS

C.-H.L. and G.G. conceived of and designed the research; C.-H.L., G.R., W.Y., M.W., W.C., B.T., J.L., K.T., M.D., A.L., N.M., M.A.L., O.R.-L.G., B.B., T.H.K., H.T., G.D. and C.A. performed experiments; C.-H.L., G.R., W.Y., O.I.L., R.P.T., F.N., N.V., P.B., Y.J.Z. and G.G. analyzed data; and C.-H.L., G.R., N.V., P.B., Y.J.Z., and G.G. wrote the paper.

COMPETING FINANCIAL INTERESTS

The authors declare competing financial interests: details are available in the [online version of the paper](#).

Reprints and permissions information is available online at <http://www.nature.com/reprints/index.html>. Publisher's note: Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

- Weiner, G.J. Building better monoclonal antibody-based therapeutics. *Nat. Rev. Cancer* **15**, 361–370 (2015).
- Nimmerjahn, F. & Ravetch, J.V. Translating basic mechanisms of IgG effector activity into next generation cancer therapies. *Cancer Immunol.* **12**, 13 (2012).
- Pincetic, A. *et al.* Type I and type II Fc receptors regulate innate and adaptive immunity. *Nat. Immunol.* **15**, 707–716 (2014).
- Li, F.J. *et al.* Emerging roles for the FCRL family members in lymphocyte biology and disease. *Curr. Top. Microbiol. Immunol.* **382**, 29–50 (2014).
- van de Donk, N.W.C.J. *et al.* Monoclonal antibodies targeting CD38 in hematological malignancies and beyond. *Immunol. Rev.* **270**, 95–112 (2016).
- Ricklin, D., Hajishengallis, G., Yang, K. & Lambris, J.D. Complement: a key system for immune surveillance and homeostasis. *Nat. Immunol.* **11**, 785–797 (2010).
- Dunkelberger, J.R. & Song, W.-C. Complement and its role in innate and adaptive immune responses. *Cell Res.* **20**, 34–50 (2010).
- Hess, C. & Kemper, C. Complement-mediated regulation of metabolism and basic cellular processes. *Immunity* **45**, 240–254 (2016).
- Sondermann, P. & Szymkowski, D.E. Harnessing Fc receptor biology in the design of therapeutic antibodies. *Curr. Opin. Immunol.* **40**, 78–87 (2016).
- Bruhns, P. & Jönsson, F. Mouse and human FcR effector functions. *Immunol. Rev.* **268**, 25–51 (2015).
- Taylor, R.P. & Lindorfer, M.A. The role of complement in mAb-based therapies of cancer. *Methods* **65**, 18–27 (2014).
- Di Gaetano, N. *et al.* Complement activation determines the therapeutic activity of rituximab *in vivo*. *J. Immunol.* **171**, 1581–1587 (2003).
- Bruhns, P. *et al.* Specificity and affinity of human Fcγ receptors and their polymorphic variants for human IgG subclasses. *Blood* **113**, 3716–3725 (2009).
- Lux, A., Yu, X., Scanlan, C.N. & Nimmerjahn, F. Impact of immune complex size and glycosylation on IgG binding to human FcγRs. *J. Immunol.* **190**, 4315–4323 (2013).
- Golay, J. & Introna, M. Mechanism of action of therapeutic monoclonal antibodies: promises and pitfalls of *in vitro* and *in vivo* assays. *Arch. Biochem. Biophys.* **526**, 146–153 (2012).
- Vafa, O. *et al.* An engineered Fc variant of an IgG eliminates all immune effector functions via structural perturbations. *Methods* **65**, 114–126 (2014).
- Arduin, E. *et al.* Highly reduced binding to high and low affinity mouse Fcγ receptors by L234A/L235A and N297A Fc mutations engineered into mouse IgG2a. *Mol. Immunol.* **63**, 456–463 (2015).
- Schreiber, A.D. & Frank, M.M. Role of antibody and complement in the immune clearance and destruction of erythrocytes. II. Molecular nature of IgG and IgM complement-fixing sites and effects of their interaction with serum. *J. Clin. Invest.* **51**, 583–589 (1972).
- Ackerman, M. & Nimmerjahn, F. *Antibody Fc: Linking Adaptive and Innate Immunity* (Academic Press, 2013).
- Wang, S.-Y., Racila, E., Taylor, R.P. & Weiner, G.J. NK-cell activation and antibody-dependent cellular cytotoxicity induced by rituximab-coated target cells is inhibited by the C3b component of complement. *Blood* **111**, 1456–1463 (2008).
- Harvey, B.R. *et al.* Anchored periplasmic expression, a versatile technology for the isolation of high-affinity antibodies from *Escherichia coli*-expressed libraries. *Proc. Natl. Acad. Sci. USA* **101**, 9193–9198 (2004).
- Jung, S.T. *et al.* Effective phagocytosis of low Her2 tumor cell lines with engineered, aglycosylated IgG displaying high FcγRIIa affinity and selectivity. *ACS Chem. Biol.* **8**, 368–375 (2013).

23. Borrok, M.J., Jung, S.T., Kang, T.H., Monzingo, A.F. & Georgiou, G. Revisiting the role of glycosylation in the structure of human IgG Fc. *ACS Chem. Biol.* **7**, 1596–1602 (2012).
24. Pawluczko, A.W. *et al.* Binding of submaximal C1q promotes complement-dependent cytotoxicity (CDC) of B cells opsonized with anti-CD20 mAbs ofatumumab (OFA) or rituximab (RTX): considerably higher levels of CDC are induced by OFA than by RTX. *J. Immunol.* **183**, 749–758 (2009).
25. Teeling, J.L. *et al.* Characterization of new human CD20 monoclonal antibodies with potent cytolytic activity against non-Hodgkin lymphomas. *Blood* **104**, 1793–1800 (2004).
26. Boross, P. & Leusen, J.H.W. Mechanisms of action of CD20 antibodies. *Am. J. Cancer Res.* **2**, 676–690 (2012).
27. Teeling, J.L. *et al.* The biological activity of human CD20 monoclonal antibodies is linked to unique epitopes on CD20. *J. Immunol.* **177**, 362–371 (2006).
28. Diebolder, C.A. *et al.* Complement is activated by IgG hexamers assembled at the cell surface. *Science* **343**, 1260–1263 (2014).
29. Beum, P.V. *et al.* Complement activation on B lymphocytes opsonized with rituximab or ofatumumab produces substantial changes in membrane structure preceding cell lysis. *J. Immunol.* **181**, 822–832 (2008).
30. Gorter, A. & Meri, S. Immune evasion of tumor cells using membrane-bound complement regulatory proteins. *Immunol. Today* **20**, 576–582 (1999).
31. Min, X. *et al.* Expression and regulation of complement receptors by human natural killer cells. *Immunobiology* **219**, 671–679 (2014).
32. Ramos, O.F., Sármay, G., Klein, E., Yefenof, E. & Gergely, J. Complement-dependent cellular cytotoxicity: lymphoblastoid lines that activate complement component 3 (C3) and express C3 receptors have increased sensitivity to lymphocyte-mediated lysis in the presence of fresh human serum. *Proc. Natl. Acad. Sci. USA* **82**, 5470–5474 (1985).
33. Romain, G. *et al.* Antibody Fc engineering improves frequency and promotes kinetic boosting of serial killing mediated by NK cells. *Blood* **124**, 3241–3249 (2014).
34. Bologna, L. *et al.* Ofatumumab is more efficient than rituximab in lysing B chronic lymphocytic leukemia cells in whole blood and in combination with chemotherapy. *J. Immunol.* **190**, 231–239 (2013).
35. DiLillo, D.J. & Ravetch, J.V. Differential Fc-receptor engagement drives an anti-tumor vaccinal effect. *Cell* **161**, 1035–1045 (2015).
36. Klaus, G.G., Pepys, M.B., Kitajima, K. & Askonas, B.A. Activation of mouse complement by different classes of mouse antibody. *Immunology* **38**, 687–695 (1979).
37. Neuberger, M.S. & Rajewsky, K. Activation of mouse complement by monoclonal mouse antibodies. *Eur. J. Immunol.* **11**, 1012–1016 (1981).
38. Bruhns, P. Properties of mouse and human IgG receptors and their contribution to disease models. *Blood* **119**, 5640–5649 (2012).
39. Lim, S.H. *et al.* Fc gamma receptor IIb on target B cells promotes rituximab internalization and reduces clinical efficacy. *Blood* **118**, 2530–2540 (2011).
40. Vaughan, A.T. *et al.* Activatory and inhibitory Fcγ receptors augment rituximab-mediated internalisation of CD20 independent of signalling via the cytoplasmic domain. *J. Biol. Chem.* jbc.M114.593806 (2015).
41. Frank, M., Walker, R.C., Lanzilotta, W.N., Prestegard, J.H. & Barb, A.W. Immunoglobulin G1 Fc domain motions: implications for Fc engineering. *J. Mol. Biol.* **426**, 1799–1811 (2014).
42. Krissinel, E. & Henrick, K. Inference of macromolecular assemblies from crystalline state. *J. Mol. Biol.* **372**, 774–797 (2007).
43. Gaboriaud, C. *et al.* The crystal structure of the globular head of complement protein C1q provides a basis for its versatile recognition properties. *J. Biol. Chem.* **278**, 46974–46982 (2003).
44. Duncan, A.R. & Winter, G. The binding site for C1q on IgG. *Nature* **332**, 738–740 (1988).
45. Idusogie, E.E. *et al.* Mapping of the C1q binding site on rituxan, a chimeric antibody with a human IgG1 Fc. *J. Immunol.* **164**, 4178–4184 (2000).
46. Schneider, S. & Zacharias, M. Atomic resolution model of the antibody Fc interaction with the complement C1q component. *Mol. Immunol.* **51**, 66–72 (2012).
47. Krapp, S., Mimura, Y., Jefferis, R., Huber, R. & Sondermann, P. Structural analysis of human IgG-Fc glycoforms reveals a correlation between glycosylation and structural integrity. *J. Mol. Biol.* **325**, 979–989 (2003).
48. Lu, J. *et al.* Structure of FcγRI in complex with Fc reveals the importance of glycan recognition for high-affinity IgG binding. *Proc. Natl. Acad. Sci. USA* **112**, 833–838 (2015).
49. Kennedy, A.D. *et al.* Rituximab infusion promotes rapid complement depletion and acute CD20 loss in chronic lymphocytic leukemia. *J. Immunol.* **172**, 3280–3288 (2004).
50. Cartron, G. *et al.* Therapeutic activity of humanized anti-CD20 monoclonal antibody and polymorphism in IgG Fc receptor FcγRIIIa gene. *Blood* **99**, 754–758 (2002).

ONLINE METHODS

Cells and reagents. All cancer cell lines were tested for mycoplasma contamination before cytotoxicity assays. Burkitt's lymphoma Raji (ATCC CCL-86), Ramos (ATCC CRL-1596), Daudi (ATCC CCL-213) and DB cancer cells (human B cell lymphoma, ATCC CRL2289) were obtained from American Type Culture Collection. TMD8 cells (human ABC-DLBCL cell line) and HBL-1 cells (human diffuse large B-cell lymphoma cell line) were obtained from P. Tucker (University of Texas at Austin). Patient-derived primary acute lymphocytic leukemia cells were obtained from D. Lee (MD Anderson Cancer Center). All cancer cells were cultured in complete RPMI with 10% FBS (FBS). The collection of CHO cells expressing human and mouse FcγRs was described previously^{13,14,51}.

Human serum depleted of C1q or C9 was purchased from CompTech. Human FcγR and antibody proteins were produced in Expi293 cells (Invitrogen) as described previously²². All primers were synthesized by IDT (Supplementary Table 1).

Library construction and screening. IgG polypeptides were displayed on the inner membrane of *Escherichia coli* (*E. coli*) as described previously²². Random mutations were introduced on the Fc domain by three different methods (methods details, Supplementary Fig. 1b). As described previously²², the libraries were made into spheroplasts after induction with 1 mM of isopropyl-1-thio- β -D-galactopyranoside (IPTG) and 2% D-arabinose, and were labeled with 10 nM C1q-PE in the presence of 1 μ M of FcγRs as competitors and screened on a FACSAria (BD Biosciences) for seven rounds. 23 randomly selected clones were analyzed by flow cytometry using 10 nM C1q-PE or 10 nM of a tetramer of FcγRIIIa_{V158} (human FcγRIIIa of an allotype of higher affinity) and SA-PE.

Detection of lipopolysaccharide on spheroplasts of *E. coli*. Antibody M18 to PA domain 4 (10 μ g/ml) was incubated with cells in spheroplasts expressing PA domain 4. In this manner, the spheroplasts are coated with full-length glycosylated IgG to enable monitoring of the binding of C1q or FcγRs onto cells in spheroplasts. Antibody-expressing cells in spheroplasts were incubated with 10 nM C1q-PE for 1 h in PBS, pH 7.4, or with high-salt phosphate buffer (50 mM phosphate and 330 mM NaCl, pH 7.4). Binding intensity of C1q-PE was detected by flow cytometry.

Preparation of antibodies, C1q and FcγR. Genes encoding 801-Fc and 802-Fc were cloned in-frame into the mammalian expression vector pcDNA3.4 with (or without) Fab of Rituximab or ofatumumab using Gibson Assembly cloning (NEB)⁵². Expression plasmids encoding histidine-tagged mouse FcγRI-IV (His-mFcγRI-IV) were constructed by cloning gene blocks of the respective genes synthesized by IDT into pcDNA3.4 vector (Invitrogen) using HindIII and Xho I sites. IgG proteins, human and mouse FcγRs were produced as described previously²². Human C1q protein was purchased from CompTech and was labeled with R-phycoerythrin (R-PE) using an EasyLink R-PE Conjugation Kit (Abcam) according to the manufacturer's instructions.

Liquid chromatography–mass spectrometry of 801-Fc. The 801-Fc proteins were analyzed by liquid chromatography–mass spectrometry on an Orbitrap Fusion mass spectrometer (ThermoFisher). A linear gradient of 0.1% formic acid in water and 0.1% formic acid in acetonitrile over 10 min was used to elute the intact proteins from an OPTI-TRAPTM protein microtrap (Optimize Technologies). The Orbitrap Fusion was operated in Intact Protein Mode at 60,000 resolution from 400–2,000 *m/z*. Following acquisition, the data were deconvoluted using MagTran software.

Tandem mass spectrometry of crosslinked A801-Fc. A801-Fc (10 μ M) and EDC (1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride) (1 mM) were mixed in a buffer containing 20 mM MES, pH 6.0, and 50 mM NaCl. Samples were incubated for 1 d at 4 °C before being separated by SDS-PAGE. The dimer bands on the SDS-PAGE gel were cut and trypsinized, and tryptic digests were analyzed by liquid chromatography–tandem mass spectrometry on an Orbitrap Elite (Thermo Fischer). Published methods were used for analysis of data obtained by mass spectrometry and tandem mass spectrometry⁵³ and assignment of peptide spectral matches. Crosslinked peptides were

identified using Protein Metrics Byonic software custom crosslinking with EDC on glutamic acid, aspartic acid and lysine residues.

Surface-plasmon-resonance analysis. To determine the affinities of antibody variants with C1q, FcγRs and FcRn, a Biacore 3000 instrument (GE Healthcare) was used with HBS-EP running buffer (GE Healthcare). Bovine serum albumin (BSA) was immobilized in the reference channel of the CM5 sensor chip, and each antibody was immobilized on the CM5 sensor chip by amine coupling at a pH of 5.0. All FcγRs (400 nM) or the serially diluted C1q proteins (1–40 nM) were injected into the CM5 chip at 30 μ l/min for 2 min, followed by 10 min of dissociation. The chip was regenerated after each binding event with 10 mM glycine, pH 3.0, with a contact time of 1 min. FcRn- β 2-microglobulin-glutathione S-transferase (730 nM) was injected in the CM5 chip at 30 μ l/min in HBS-EP, pH 7.4, for 90 s, and the chip was regenerated after each binding event with 10 mM glycine, pH 3.0, with a contact time of 1 min. Monomeric FcRn- β 2-microglobulin (50–400 nM) was injected into the CM5 chip at 30 μ l/min in PBS, pH 6.0, for 90 s and the chip was regenerated after each binding event with 10 mM glycine, pH 3.0, with a contact time of 1 min. The resulting sensorgrams were fit with a global two-state binding model for C1q, and with a 1:1 Langmuir isotherm model for monomeric FcγRI and FcRn using Biaevaluation 3.0 software. Because C1q is hexameric protein, the measured K_D values do not correspond to the true equilibrium dissociation constants for the interaction of the C1q globular head with IgG and are used here only as a measure of the relative affinity of the C1q for each antibody.

C3b deposition assays. Cells cultured in complete RPMI-1640 medium were mixed with an equal volume of normal human serum (NHS), and then mAbs were added to a final concentration of 10 μ g/ml. After incubation for 30 min at 37 °C, the cells were washed twice with 1% BSA in PBS and developed with FITC-conjugated mouse mAb 3E7 to C3b (BioLegend; 1:50 dilution). Flow cytometry was performed on a FACSCalibur flow cytometer (BD Biosciences), and mean fluorescence intensity was converted into molecules of equivalent soluble fluorochrome with calibrated beads (Spherotech).

Solution-phase complement-activation assay. Complement activation in the absence of tumor cells was determined by measurement of the concentration of C4d, a product of the classical complement activation pathway, after incubation of 100 μ g antibody in 1 ml 90% NHS for 1 h at 37 °C. C4d concentrations were measured by enzyme-linked immunosorbent assay (MicroVue C4d EIA kit, Quidel Corporation) according to the manufacturer's instructions.

Size-exclusion chromatography (SEC). SEC was performed on Agilent 1100 high-performance liquid chromatography system using a SuperdexTM200 10/300GC, (GE Healthcare), with a mobile phase of PBS, pH 7.4, at a flow rate of 0.5 ml/min. Proteins were detected by monitoring the absorbance at 280 nm. The injection amount was 100 μ g protein in a volume of 100 μ l.

Enzyme-linked immunosorbent assay. To measure binding of antibodies to FcγRs and C1q, 96-well plates (Qiagen) were coated with 1 μ g/well of each antibody at 4 °C for 16 h and were washed three times with PBS containing 0.05% Tween20 (PBST). The plates were blocked for 1 h at 25 °C with 3% skim milk in PBS and were washed three times with PBST. FcγRs (50 nM or 500 nM) was added in plates. After 1 h of incubation at 25 °C, the plates were washed with PBST and incubated with 50 μ l of PBS containing 1:5,000 goat anti-histidine (Abcam; Cat# ab1269) or antibody to glutathione S-transferase IgG conjugated to HRP (Rockland; Cat# 600-103-200) for 1 h. After washing three times with PBST, 50 μ l TMB substrate was added per well (Thermo Scientific), 50 μ l of 1 M H₂SO₄ was added for neutralization, and the absorbance at 450 nm was recorded.

Crystallization and data collection. Crystallization conditions for the 801-Fc domain were screened initially by sparse-matrix solutions using sitting drop-vapor diffusion on Phenix instrument (Art Robins Instruments). Crystals for A801-Fc were obtained by mixing protein at a concentration of 20 mg/ml with buffer containing 50 mM sodium citrate, pH 5.0, as a pH buffer, and 0.44 M NaCl with 24% wt/vol PEG 4000 as precipitant at 1:1 ratio at 25 °C. G801-Fc at

a concentration of 10 mg/ml crystallized in 40 mM potassium phosphate, 16% wt/vol PEG 8000 and 20% vol/vol glycerol. The crystals were cryo-protected in 20% glycerol before vitrified in liquid nitrogen for data collection⁵⁴.

Crystal diffraction data were collected at the Advanced Light Source beam-line BL5.0.3 (Berkeley, CA) and Advanced Photo Source BL 23-IDD (Chicago, IL). The data were processed using the program HKL2000 (ref. 54). The diffractions were scaled to 2.3 Å for A801-Fc and 3.2 Å for G801-Fc. The statistics for data collection are summarized in **Table 3**.

Structure determination, refinement and analysis. The structures of A801 was determined by molecular replacement using wild-type Fc as the search model (PDB code: 3AVE). The first round of molecular refinement using program Phaser in CCP4 suite^{55,56} identified two dimers per asymmetric unit and revealed that the Cγ2-Cγ3 dihedral angle varied considerably from those in all other Fc structures in the database. Therefore, the Cγ2 and Cγ3 were searched separately by Phaser^{55,56}, and the initial solution was identified and then confirmed with electron-density-fitting the model. Rigid-body and CNS refinement were carried out in Phenix.refine⁵⁷. The model was optimized by iterative cycles of manual model building using COOT⁵⁸ followed by refinement with Phenix.refine⁵⁷ to improve the quality of the model. Prior to the refinement, 5% of the diffraction data were reserved as an unbiased test set for cross-validation (R_{free})⁵⁹. The final model exhibits an R_{work} of 20% and an R_{free} of 24%.

The structure of G801-Fc was also determined by molecular replacement using the structure of glycosylated human Fc as the search model (PDB code: 3AVE). A total of four Fc molecules were found per asymmetric unit. The fitted model was rebuilt using COOT⁵⁸, and several iterative cycles of optimization were carried out with manual building and refinement. The final model showed an R_{work} of 25% and an R_{free} of 29%.

Both structures were evaluated by MolProbity⁶⁰ and Procheck. For A801, the MolProbity Score was 1.71 and placed this at the 99th percentile of all structures for the similar resolution range in database. For G801-Fc, the MolProbity Score was 1.47 and reached 100th percentile. Detailed refinement statistics for A801-Fc and G801-Fc are summarized in **Table 3**. Figures were prepared using PyMol (DeLano, 2002)⁶¹.

C1q cell-surface-binding assay. Raji cells were resuspended in 5% NHS and then incubated with mAbs at a final concentration of 10 μg/ml at 37 °C for various times (0–30 min). Classical complement activation reactions were then quenched with 20 volumes of ice-cold 1% BSA-PBS. The cells were pelleted and washed once and then were probed with FITC anti-C1q (Dako; Cat# F0254; 1:200 dilution) for 30 min at 25 °C. The samples were washed and then analyzed by flow cytometry (FACSCalibur flow cytometer; BD Biosciences), and mean fluorescence intensity was converted to molecules of equivalent soluble fluorochrome with calibrated beads (Spherotech).

Binding of ICs to FcγRs expressed on CHO cells. ICs were generated by co-incubation of 5 μg/ml of variants of anti-TNP IgG (clone 7B4; provided by F.N.) and 2.5 μg/ml of TNP-coupled BSA (TNP₄-BSA or TNP₃₂-BSA; Biosearch Technologies) for 3 h at 25 °C (ref. 14). The relative size of TNP₄-BSA and TNP₃₂-BSA ICs was analyzed by polyethylene glycol (PEG) precipitation. ICs were then incubated with 1 × 10⁵ CHO cells stably expressing human FcγRs for 1 h with gentle shaking at 4 °C, followed by detection of bound ICs by flow cytometry using a PE-conjugated goat anti-human IgG F(ab')₂ fragment at 0.5 mg/ml (Jackson Laboratories; Cat# 109-096-006). ICs were also generated by mixing 10 μg/ml of antibodies and 5 μg/ml of PE-conjugated F(ab')₂ goat anti-human IgG F(ab')₂ (identified above) for 30 min at 37 °C¹³. CHO cells expressing mouse FcγRs were incubated with these ICs for 1 h on ice and were analyzed by flow cytometry with the method described above. Data were analyzed with Flow Cytometry Analysis Software (FlowJo).

Complement-dependent cytotoxicity (CDC) assays. CDC assays were performed as described previously⁵². In brief, serially diluted antibodies were incubated with 25% pooled human serum and cancer cells loaded with calcein AM (Life Technologies) at 37 °C for 1 h in 96-well plates. The plates were centrifuged at 1,000g for 10 min and then the supernatants were collected. The released calcein-AM was detected by spectrophotometry using an excitation

wavelength of 485 nm and emission wavelength of 535 nm. The percent of tumor cell lysis was calculated relative to SDS lysis buffer. The detailed equation is provided in **Supplementary Figure 4**. CDC assays with serum depleted of C9 or pooled mouse serum were performed exactly as described above.

PBMC or PMN purification. All *in vitro* assays were performed under a protocol approved by the UT Austin Institutional Review Board (IRB). Human peripheral blood mononuclear cells (PBMCs) and polymorphonuclear leukocytes (PMNs) were isolated from human blood from healthy donors on the day before the ADCC, CDCC and CMC assays. In brief, 50 ml of human blood was collected in heparinized vials (BD biosciences). 25 ml of blood was layered over 25 ml of Histopaque (Sigma-Aldrich) in 50-ml conical tubes. The tubes were centrifuged at 1,000g for 30 min in a swing-out bucket with no brakes. PBMCs were aspirated in the interphase between Histopaque and medium, and PMNs were collected from the pellet. Both human PBMCs and PMNs were resuspended with red-blood-cell lysis buffer (155 mM NH₄Cl, 12 mM NaHCO₃ and 0.1 mM EDTA) and were washed twice with PBS.

Complement- or FcγR-mediated cellular cytotoxicity assays. For ADCC assays, 5 × 10⁴ human PBMCs or PMNs were mixed with various concentrations of IgG variants and 5 × 10³ of calcein-AM-loaded cancer cells. For complement-mediated cell cytotoxicity assays, 25% of serum depleted of C9 or pooled human serum was added in wells containing PBMCs (or activated-PMNs), antibodies and cancer cells. Before use, PMNs were stimulated by GM-CSF (10 ng/ml) for 24 h. After 4 h at 37 °C, the fraction of lysed tumor cells was determined by the same method as CDC assay. For all assays, an E:T ratio of 10:1 was used.

Complement-receptor-inhibition assays. Calcein-AM-loaded cancer cells were pre-incubated with antibodies for 30 min. Antibody-opsonized Raji cells were incubated with effector cells (PBMC or PMN) in RPMI-1640 medium supplemented with 25% serum depleted of C1q. After 4 h at 37 °C, the fraction of lysed tumor cells was detected as described for CDC assays above. For all assays, an E:T ratio of 10:1 was used.

Blockade assays for CR3 or CR4 were also performed. 10 μg/ml of anti-CR3 or anti-CR4 (BioLegend; Cat# 301302 or Cat# 301616) was pre-incubated with effector cells for 30 min. Calcein-AM-loaded and antibody-opsonized CD20⁺ cancer cells were then incubated with anti-CR3- or anti-CR4-coated effector cells in RPMI-1640 medium supplemented with 25% serum depleted of C9. After 4 h at 37 °C, the fraction of lysed tumor cells was detected as described for CDC assays above. For all assays, an E:T ratio of 10:1 was used.

Phagocytosis assays (ADCP or CDCP). CD14⁺ monocytes were first isolated from PBMCs by EasySep (STEMCELL Inc). Monocytes were differentiated into M1 macrophages by culture for 7 d in RPMI-1640 medium containing 10% FBS and 100 ng/ml GM-CSF in 5% CO₂ incubator. On the day of the assay, adherent macrophages were detached with HyQase (GE Healthcare). For ADCP assays, 1 × 10⁵ M1 macrophages were mixed with various concentrations of IgG variants and 1 × 10⁴ of calcein-AM-loaded cancer cells. For CDCP assays, 25% serum depleted of C9 was added to wells containing M1 macrophage, antibodies, and cancer cells. After 2 h, M1 macrophages were labeled with anti-human-CD11b-APC (BioLegend; Cat# 301310) and anti-human-CD14-APC (BioLegend; Cat# 301807). Phagocytosis was evaluated by flow cytometry on an LSRFortessa (BD Bioscience), and results are reported as the ratio of cells positive for both calcein-AM and CD11b-CD14 to the total number of tumor cells in the sample.

Fluorescent images of macrophages phagocytosing Raji cells were obtained by confocal microscopy using calcein-AM-loaded Raji cells opsonized with 5 μg/ml of Rituxan, RA801 and RA802 and incubated with serum depleted of C9 and M1 macrophages at 37 °C for 2 h. Approximately 1 × 10⁶ labeled Raji cells and 1 × 10⁶ macrophages were co-incubated in a total volume of 1 ml. Subsequently, the co-incubated cells were labeled with anti-human-CD11b-APC (BioLegend; Cat# 301310) and anti-human-CD14-APC (BioLegend; Cat# 301807). Phagocytosis was visualized by confocal microscopy using a Zeiss LSM 710/Elyra S.1.

TIMING assays. Thin Nanowell arrays were fabricated following an adaptation of previously described protocol⁶², and a piece 2-cm × 2-cm in size was

cut and fixed on a #1.5 glass-bottomed, 50-mm diameter Petri dish (Ted Pella). The single-cell cytotoxicity assay was performed as described before³³. In brief, Raji cells (targets) and freshly isolated blood NK cells (effectors)³³ were labeled for 2 min with 1 μ M of red PKH26 and green PKH67 (Sigma-Aldrich), respectively. Except for the control in which Raji cells were not opsonized with antibodies, Raji cells were incubated with 1 μ g/ml of RA801 or rituximab for 30 min at 4 °C in complete medium containing serum depleted of C9. NK cells and Raji cells were loaded sequentially onto Nanowell arrays at a concentration of 1×10^6 cells/ml, allowed to settle for 3 min, and the entire chip was immersed in phenol-red-free complete medium containing 1/60 (vol/vol) annexin V-AlexaFluor-647 (Life Technologies) and, depending on the condition, the antibodies. Images were acquired on an AxioObserver mounted with a 5% CO₂, 37 °C and 100% humidity incubator, piloted with Zen software (Carl Zeiss) and fitted with a Hamamatsu EM-CCD camera and a 20 \times 0.8 NA objective. TIMING images were taken for 6 h at intervals of 5 min, and data were processed and analyzed as described previously⁶³. In brief, a semi-automated pipeline of image analysis takes care of image renaming and timelapse stacking, spectral overlap 'un-mixing', background subtraction, cropping of images around the wells and detection of cells by binarization of cell fluorescence, tracking of cell along time, and feature computation.

Animal studies. All animal experiments were performed under a protocol approved by UT Austin institutional Animal Care and Use Committee (IACUC). 1×10^6 Ramos cells in 100 μ l PBS with 50% Matrigel (BD Bioscience) were injected subcutaneously in the right flank of athymic nude mice (Jackson Laboratory). Administration of antibodies (10 mg/kg) or PBS was begun when the tumor area reached to an average of 30 mm² and was repeated a total three times on day 7, day 10 and day 13. Tumor diameters were measured every 3–4 d with a caliper, and tumor areas were calculated by the formula (length) \times (width) \times π . Mice were euthanized when the tumor size reached 2,500 mm³ in volume.

1×10^5 EL-hCD20-Luc2 cells in culture medium were injected subcutaneously into either NMRI-*Foxn1*^{null}/*Foxn1*^{null} mice ($n = 10$, Janvier Labs) or Fc γ R-null NMRI-*Foxn1*^{null}/*Foxn1*^{null} mice generated by intercross of Fc γ R-null mice (*Fcgr1*, *Fcgr2b*, *Fcgr3* and *Fcgr4*) on the 129/C57BL/6 background and NMRI-*Foxn1*^{null}/*Foxn1*^{null} mice. Antibodies (5 mg/kg) or PBS was intravenously administrated three times on days 1, 2 and 3. For measurement of tumor growth by bioluminescence (IVIS SpectrumCT, PerkinElmer), 750 μ g of D-luciferin (PerkinElmer) was intraperitoneal injected after anesthesia. Total photon flux (photons/seconds) was calculated using Living-Image-v4.5 software.

Anaphylaxis assay. Heat aggregated IgG variants, Fab as negative control, mouse Rituximab as positive control and RA801 were heat aggregated by incubation

at 63 °C for 1 h. 600 μ g of each heat-aggregated antibodies was injected into C57BL/6J mice (The Jackson Laboratory) intravenously. Mouse core body temperature was measured every 5 min using a rectal thermoprobe.

Fc γ RIIb-mediated internalization assay. CD20⁺Fc γ RIIb⁺ Raji, HBL-1 or TMD8 cells were incubated with 10 μ g of antibodies for 0, 2, 4 or 6 h in RPMI-1640 medium supplemented with 10% FBS (Invitrogen). The level of cell-surface bound antibodies was detected by goat anti-human Fc with FITC (Abcam; Cat# ab97224; 1:200 dilution).

Data analysis and statistics. Data were post-processed using Excel and GraphPad Prism. Statistical testing was run using Fisher's exact test when contingency numbers were compared, log-rank test when survival curves were compared, and elsewhere we used analysis of variance and *t*-tests.

Data availability. All the other data that support the findings of this study are available from the corresponding author upon request. PDB accession codes for the protein structural data are 5V43 for A801-Fc and 5V4E for G801-Fc.

51. Mancardi, D.A. *et al.* Fc γ RIV is a mouse IgE receptor that resembles macrophage Fc ϵ RI in humans and promotes IgE-induced lung inflammation. *J. Clin. Invest.* **118**, 3738–3750 (2008).
52. Kelton, W. *et al.* IgGA: a "cross-isotype" engineered human Fc antibody domain that displays both IgG-like and IgA-like effector functions. *Chem. Biol.* **21**, 1603–1609 (2014).
53. Lee, C.-F., Paull, T.T. & Person, M.D. Proteome-wide detection and quantitative analysis of irreversible cysteine oxidation using long column UPLC-MS/MS. *J. Proteome Res.* **12**, 4302–4315 (2013).
54. Otwinowski, Z. & Minor, W. Processing of X-ray diffraction data collected in oscillation mode. *Methods Enzymol.* **276**, 307–326 (1997).
55. Winn, M.D. *et al.* Overview of the CCP4 suite and current developments. *Acta Crystallogr. D Biol. Crystallogr.* **67**, 235–242 (2011).
56. McCoy, A.J. *et al.* Phaser crystallographic software. *J. Appl. Crystallogr.* **40**, 658–674 (2007).
57. Adams, P.D. *et al.* PHENIX: a comprehensive Python-based system for macromolecular structure solution. *Acta Crystallogr. D Biol. Crystallogr.* **66**, 213–221 (2010).
58. Emsley, P., Lohkamp, B., Scott, W.G. & Cowtan, K. Features and development of Coot. *Acta Crystallogr. D Biol. Crystallogr.* **66**, 486–501 (2010).
59. Brünger, A.T. Free R value: a novel statistical quantity for assessing the accuracy of crystal structures. *Nature* **355**, 472–475 (1992).
60. Chen, V.B. *et al.* MolProbity: all-atom structure validation for macromolecular crystallography. *Acta Crystallogr. D Biol. Crystallogr.* **66**, 12–21 (2010).
61. Terwilliger, T.C. Maximum-likelihood density modification. *Acta Crystallogr. D Biol. Crystallogr.* **56**, 965–972 (2000).
62. Liadi, I., Roszik, J., Romain, G., Cooper, L.J.N. & Varadarajan, N. Quantitative high-throughput single-cell cytotoxicity assay for T cells. *J. Vis. Exp.* **72**, e50058 (2013).
63. Merouane, A. *et al.* Automated profiling of individual cell-cell interactions from high-throughput time-lapse imaging microscopy in nanowell grids (TIMING). *Bioinformatics* **31**, 3189–3197 (2015).

Corrigendum: IgG Fc domains that bind C1q but not effector Fc γ receptors delineate the importance of complement-mediated effector functions

Chang-Han Lee , Gabrielle Romain, Wupeng Yan , Makiko Watanabe, Wissam Charab, Biliana Todorova, Jiwon Lee, Kendra Triplett, Moses Donkor, Oana I Lungu, Anja Lux, Nicholas Marshall, Margaret A Lindorfer, Odile Richard-Le Goff, Bianca Balbino, Tae Hyun Kang, Hidetaka Tanno, George Delidakis, Corrine Alford, Ronald P Taylor, Falk Nimmerjahn, Navin Varadarajan, Pierre Bruhns , Yan Jessie Zhang & George Georgiou

Nat. Immunol.; doi:10.1038/ni.3770; corrected online 27 June 2017

In the version of this article initially published online, the labels identifying each plot in Figure 1b were missing. The labels are as follows (left to right): CHO, Fc γ RI, Fc γ RIIa_{R131}, Fc γ RIIa_{H131}, Fc γ RIIb, Fc γ RIIIa_{F158} and Fc γ RIIIa_{V158}. Also, the reference cited in the accompanying legend (ref. 21) is incorrect. The correct reference is ref. 14. The errors have been corrected in the print, PDF and HTML versions of this article.